



DESERT TORTOISE COUNCIL

3807 Sierra Highway #6-4514

Acton, CA 93510

www.deserttortoise.org

eac@deserttortoise.org

Via email only

10 May 2023

Attn: Kenny Kendrick
Bureau of Land Management
4701 N. Torrey Pines Drive
Las Vegas, NV 89130
kkendrick@blm.gov, [BLM NV LVFO Muddy Mt TMP@blm.gov](mailto:BLM_NV_LVFO_Muddy_Mt_TMP@blm.gov)

RE: Scoping Comments - Muddy Mountains Travel Management Plan (DOI-BLM-NV-S010-2023-0040-EA)

Dear Mr. Kendrick,

The Desert Tortoise Council (Council) is a non-profit organization comprised of hundreds of professionals and laypersons who share a common concern for wild desert tortoises and a commitment to advancing the public's understanding of desert tortoise species. Established in 1975 to promote conservation of tortoises in the deserts of the southwestern United States and Mexico, the Council routinely provides information and other forms of assistance to individuals, organizations, and regulatory agencies on matters potentially affecting desert tortoises within their geographic ranges.

Both our physical and email addresses are provided above in our letterhead for your use when providing future correspondence to us. When given a choice, we prefer that you email to us future correspondence, as mail delivered via the U.S. Postal Service may take several days to be delivered. Email is an "environmentally friendlier way" of receiving correspondence and documents rather than "snail mail."

We appreciate that the Bureau of Land Management (BLM) contacted the Council directly via email on 4/12/2023 for the opportunity to provide scoping comments on the above-referenced project. Given the location of the proposed project in habitats likely occupied by Mojave desert tortoise (*Gopherus agassizii*) (synonymous with Agassiz's desert tortoise), our comments pertain to enhancing protection of this species during activities funded, authorized, or carried out by the BLM, which we assume will be added to the Decision Record for this project as needed. Please accept, carefully review, and include in the relevant project file the Council's following comments and attachments for the proposed project.

The Mojave desert tortoise is among the top 50 species on the list of the world's most endangered tortoises and freshwater turtles. The International Union for Conservation of Nature's (IUCN) Species Survival Commission, Tortoise and Freshwater Turtle Specialist Group, now considers the Mojave desert tortoise to be Critically Endangered (Berry et al. 2021) "... based on population reduction (decreasing density), habitat loss of over 80% over three generations (90 years), including past reductions and predicted future declines, as well as the effects of disease (upper respiratory tract disease/mycoplasmosis). *Gopherus agassizii* (*sensu stricto*) comprises tortoises in the most well-studied 30% of the larger range; this portion of the original range has seen the most human impacts and is where the largest past population losses had been documented. A recent rigorous rangewide population reassessment of *G. agassizii* (*sensu stricto*) has demonstrated continued adult population and density declines of about 90% over three generations (two in the past and one ongoing) in four of the five *G. agassizii* recovery units and inadequate recruitment with decreasing percentages of juveniles in all five recovery units."

This status, in part, prompted the Council to join Defenders of Wildlife and Desert Tortoise Preserve Committee (Defenders of Wildlife et al. 2020) to petition the California Fish and Game Commission in March 2020 to elevate the listing of the Mojave desert tortoise from threatened to endangered in California.

The following project description is given in BLM's public notice: "The Muddy Mountains planning area is approximately 123,400 acres of Bureau of Land Management (BLM) public lands located approximately 50 miles northeast of Las Vegas and approximately 25 miles southwest of Logandale in Southern Nevada. The adjacent lands are managed by several land management entities: the National Park Service, Nevada State Parks and Tribal lands. The planning area also includes and is bounded by private lands. Muddy Mountains contains unique land features, such as colorful sandstone outcrops, rugged limestone cliffs, and canyons. It is also home to sensitive cultural and natural resources that require protection and management.

"The planning area provides numerous recreational opportunities and is a destination for many types of off-highway-vehicle (OHV) uses. Muddy Mountains was designated as a Special Recreation Management Area (SRMA) in the 1998 Las Vegas Resource Management Plan (RMP). The BLM Las Vegas Field Office (LVFO) is preparing a Travel Management Plan (TMP) and Environmental Assessment (EA) for the Muddy Mountains Travel Management Area (TMA). The TMA decision area consists of only BLM-managed lands. The 1998 Las Vegas RMP management direction states that the objective of the Muddy Mountains area is 'to provide semi-primitive recreation opportunities and integrated management of wildlife habitat, cultural resources, and other recreational uses.' The TMP will identify, and designate routes open to the public, routes for administrative use only, and routes to be closed and rehabilitated, as well as determine the appropriate management for those routes. The TMP may also identify new routes to help meet the goals and objectives of the 1998 Las Vegas RMP, and amendments, as well as other applicable laws, regulations, and policies including, but not limited to, the Federal Lands Policy and Management Act (FLPMA) and the National Environmental Policy Act (NEPA)."

We have expressed in numerous comment letters to the Las Vegas office of the BLM our concern with making decisions based on the outdated, obsolete 1998 Las Vegas Resource Management Plan (Las Vegas RMP; BLM 1998a, 1998b). In fact, it is our understanding that, if not already, the BLM will soon be revising the Las Vegas RMP, which we fully support. We know from multiple documents published since 1998 (Allison and McLuckie 2018, USFWS 2014a, 2014b, 2015, 2016, 2017, 2018, 2019, 2020, 2022a, 2022b) that the plight of the tortoise on lands mostly managed by BLM have declined substantially throughout the listed range of the Mojave desert, which was not a consideration in the 1998 formulation of the RMP. Please be sure that the Draft Environmental Assessment (DEA) for this TMP adequately summarizes the current status of the desert tortoise in the region. To facilitate this request, we provide Appendix A, which summarizes the current status of tortoises throughout the listed range including the Northeastern Recovery Unit, which is where most or all of the planning area/proposed action appears to be located.

The presence of roads even with low vehicle use has multiple adverse effects on the desert tortoise and its habitats. These include the deterioration/loss of wildlife habitat, hydrology, geomorphology, and air quality; increased competition and predation (including by humans); and the loss of naturalness or pristine qualities, all of which must be analyzed in the DEA. We ask that the DEA fully divulge and assess these and other impacts associated with recreational vehicle use. To facilitate this request, we herein provide BLM with Appendix B, which is a partial bibliography of impacts associated with recreational vehicle use on arid lands. We expect the authors of the DEA to familiarize themselves with this literature to better understand direct, indirect, and cumulative impacts so that they will analyze the direct, indirect, and cumulative impacts of the proposed action to the tortoise and its habitat, including habitat for connectivity and make informed management decisions to curtail these impacts before they occur.

Please include in the DEA analyses the five major categories of primary road effects to the tortoise and special status species: (1) wildlife mortality from collisions with vehicles; (2) hindrance/barrier to animal movements thereby reducing access to resources and mates; (3) degradation of habitat quality; (4) habitat loss caused by disturbance effects in the wider environment and from the physical occupation of land by the road; and (5) subdividing animal populations into smaller and more vulnerable fractions (Jaeger et al. 2005a, 2005b, Roedenbeck et al. 2007).

We note in the BLM's "Process and Schedule" graphic that "Route Inventory" is supposed to be initiated (or finalized, which is not clear) in December 2022, although we did not see any such document(s) on the eplanning website. There is a map with the TMP boundary but no routes are depicted. Please be sure that the DEA explains how this route inventory was (is being) derived and if it has (or will be) ground-truthed. It has been our experience that washes often appear on satellite imagery as routes when in fact they are natural features. Although we strongly recommend that the final proposed route inventory be fully ground-truthed, if that is not possible, we ask that the proposed network be superimposed over U.S. Geological Survey (USGS) maps to see if any of the future open routes are located in washes. Given the importance of washes to tortoises (Jennings and Berry 2015), we ask that BLM avoid designating open routes in washes. Further, we request that maps of the open, closed, and limited routes be displayed in maps in the DEA for each alternative.

Has BLM completed an inventory of tortoise abundance within the TMP planning area? If field data are unavailable, we ask that BLM use available models (e.g., Nussear 2009, Gray et al. 2019, etc.) to which the proposed route network can be compared. We ask that BLM be conscientious about minimizing the number of routes in those portions of the TMP that are known to contain higher densities of tortoises, either based on field studies (preferred) or models (less preferred). The DEA should use the best available science in analyzing impacts of the proposed action to the tortoise/tortoise habitat and selecting open and limited routes. For example, the DEA should include results and management recommendations from modeling threats to the tortoise conducted by Tuma et al. (2016) in the nearby Gold Butte-Pakoon Tortoise Conservation Area in its analyses of these impacts (i.e., threats causing habitat degradation and slight increases in tortoise mortality over a broad area, such as livestock grazing and human presence (e.g., OHV use) are greater contributors to tortoise population declines than patchily distributed threats.

The DEA should include appropriate mitigation and monitoring plans for all direct, indirect, and cumulative effects to the tortoise and its habitats; the mitigation and monitoring plan should use the best available science with a commitment to implement the mitigation commensurate to impacts to the tortoise and its habitats. Mitigation and monitoring should include a fully-developed raven management plan; weed management plan; fire management plan; compensation plan for the degradation and loss of tortoise habitat that includes protection of the acquired, improved, and restored habitat in perpetuity for the tortoise from future development and human use; a plan to develop and implement an effective education program; a plan for effective law enforcement to prevent route proliferation; and habitat restoration plan for closed routes. These plans should be part of the DEA so that the public has an opportunity to provide input on the various plans. If the plans cannot be developed in time to accompany the DEA, the BLM should commit to providing the draft plans to affected interests, including the Council, when they become available, prior to finalizing them.

These mitigation and monitoring plans should include an implementation schedule that is tied to key actions associated with route designation, signing open routes, and restoration phases of the project so that mitigation occurs concurrently with or in advance of the impacts. The Council has found that installing bright red Carsonite signs on closed routes is not effective; in fact, some routes that would not have otherwise been obvious are subject to use because of these signs. The Council prefers that closed routes be physically eradicated using vertical mulching and other techniques that eliminate the routes, and that the TMP has a schedule for closing routes. We also ask that the BLM prioritize the closure of routes based on tortoise densities derived from field studies or modelling. The plans should specify success criteria, include a monitoring plan to collect data to determine whether success criteria have been met, and identify actions that would be required if the mitigation measures do not meet the success criteria. Specific remedial measures, such as increased law enforcement, enhanced education, closure of problematic routes, etc. should be included in the plan.

Given these observations concerning closing routes, the Council provides Appendix C as a partial list of literature that will help inform the BLM in implementing restoration techniques for closed routes. In 2016, the Council funded a document (Abella and Berry 2016) presenting best management practices for arid restoration techniques, which is included in the link in the footnote¹. Also, we suggest that BLM use the attached document (Appendix D) by Abella et al. (2023) and listed in Appendix C. Habitat Restoration Bibliography for the Mojave Desert.

¹ <https://www.dropbox.com/s/nx1b5m2b5ehya12/%23Abella%20and%20Berry%202016.pdf?dl=0>

Given their impacts on tortoises, the DEA must analyze if the TMP would result in an increase of common ravens and other predators of the desert tortoise in the planning area. Future operations must include provisions for monitoring and managing raven predation on tortoises as a result of the proposed action. The monitoring and management plan must include reducing human subsidies for food, water, and sites for nesting, roosting, and perching to address local impacts.

With regards to cumulative effects, the DEA must list and discuss all project impacts within the region including future state, federal, and private actions affecting listed species on state, federal, and private lands. In particular, we ask that the DEA provide a thorough analysis of the loss and degradation of tortoises and occupied habitats throughout Nevada, particularly in the southern part of the state.

Please see *Grand Canyon Trust v. F.A.A.*, 290 F.3d 339, 345-46 (D.C. Cir. 2002) in which the court decided that federal agencies must analyze the cumulative impacts of actions in environmental assessments. In the cumulative effects analysis of the DEA, please ensure that the CEQ's "Considering Cumulative Effects under the National Environmental Policy Act" (1997) is followed, including the eight principles, when analyzing cumulative effects of the proposed action to the tortoise and its habitats. CEQ states, "Determining the cumulative environmental consequences of an action requires delineating the cause-and-effect relationships between the multiple actions and the resources, ecosystems, and human communities of concern. The range of actions that must be considered includes not only the project proposal but all connected and similar actions that could contribute to cumulative effects." The analysis "must describe the response of the resource to this environmental change." Cumulative impact analysis should "address the sustainability of resources, ecosystems, and human communities."

CEQ's guidance on how to analyze cumulative environmental consequences, which contains eight principles listed below:

1. Cumulative effects are caused by the aggregate of past, present, and reasonable future actions.

The effects of a proposed action on a given resource, ecosystem, and human community, include the present and future effects added to the effects that have taken place in the past. Such cumulative effects must also be added to the effects (past, present, and future) caused by all other actions that affect the same resource.

2. Cumulative effects are the total effect, including both direct and indirect effects, on a given resource, ecosystem, and human community of all actions taken, no matter who (federal, non-federal, or private) has taken the actions.

Individual effects from disparate activities may add up or interact to cause additional effects not apparent when looking at the individual effect at one time. The additional effects contributed by actions unrelated to the proposed action must be included in the analysis of cumulative effects.

3. Cumulative effects need to be analyzed in terms of the specific resource, ecosystem, and human community being affected.

Environmental effects are often evaluated from the perspective of the proposed action. Analyzing cumulative effects requires focusing on the resources, ecosystem, and human community that may be affected and developing an adequate understanding of how the resources are susceptible to effects.

4. It is not practical to analyze the cumulative effects of an action on the universe; the list of environmental effects must focus on those that are truly meaningful.

For cumulative effects analysis to help the decision maker and inform interested parties, it must be limited through scoping to effects that can be evaluated meaningfully. The boundaries for evaluating cumulative effects should be expanded to the point at which the resource is no longer affected significantly or the effects are no longer of interest to the affected parties.

5. Cumulative effects on a given resource, ecosystem, and human community are rarely aligned with political or administrative boundaries.

Resources are typically demarcated according to agency responsibilities, county lines, grazing allotments, or other administrative boundaries. Because natural and sociocultural resources are not usually so aligned, each political entity actually manages only a piece of the affected resource or ecosystem. Cumulative effects analysis on natural systems must use natural ecological boundaries and analysis of human communities must use actual sociocultural boundaries to ensure including all effects.

6. Cumulative effects may result from the accumulation of similar effects or the synergistic interaction of different effects.

Repeated actions may cause effects to build up through simple addition (more and more of the same type of effect), and the same or different actions may produce effects that interact to produce cumulative effects greater than the sum of the effects.

7. Cumulative effects may last for many years beyond the life of the action that caused the effects.

Some actions cause damage lasting far longer than the life of the action itself (e.g., acid mine damage, radioactive waste contamination, species extinctions). Cumulative effects analysis needs to apply the best science and forecasting techniques to assess potential catastrophic consequences in the future.

8. Each affected resource, ecosystem, and human community must be analyzed in terms of its capacity to accommodate additional effects, based on its own time and space parameters.

Analysts tend to think in terms of how the resources, ecosystem, and human community will be modified given the action's development needs. The most effective cumulative effects analysis focuses on what is needed to ensure long-term productivity or sustainability of each resource impacted by the proposed action including the Mojave desert tortoise.

We appreciate this opportunity to provide comments on this project and trust they will help protect tortoises during any resulting authorized activities. Herein, we reiterate that the Desert Tortoise Council wants to be identified as an Affected Interest for this and all other projects funded, authorized, or carried out by the BLM that may affect species of desert tortoises, and that any subsequent environmental documentation for this project is provided to us at the contact information listed above. Additionally, we ask that you respond in an email that you have received this comment letter so we can be sure our concerns have been registered with the appropriate personnel and office for this project.

Respectfully,



Edward L. LaRue, Jr., M.S.
Ecosystems Advisory Committee, Chairperson
Desert Tortoise Council

cc. Glen Knowles, Field Supervisor, Southern Nevada Field Office (Las Vegas), U.S. Fish and Wildlife Service, glen_knowles@fws.gov
Jon Raby, Nevada State Director, Bureau of Land Management, jraby@blm.gov

Attachments:

Appendix A. Demographic Status and Trend of the Mojave Desert Tortoise (*Gopherus agassizii*)
Appendix B. Bibliography on road impacts in desert ecosystems
Appendix C. Habitat Restoration Bibliography for the Mojave Desert
Appendix D. Techniques for Restoring Damaged Mojave and Western Sonoran Habitats, Including Those for Threatened Desert Tortoises and Joshua Trees.

Literature Cited

Abella, S.A., K.H. Berry, and S. Ferrazzano. 2023. Techniques for Restoring Damaged Mojave and Western Sonoran Habitats, Including Those for Threatened Desert Tortoises and Joshua Trees. *Desert Plants Journal* Volume 38, Number 2, April 2023, pages 4-52.

Allison L.J. and A.M. McLuckie. 2018. Population trends in Mojave desert tortoises (*Gopherus agassizii*). *Herpetological Conservation and Biology*. 2018 Aug 1;13(2):433-52. http://www.herpconbio.org/Volume_13/Issue_2/Allison_McLuckie_2018.pdf or <https://www.fws.gov/media/allison-and-mcluckie2018mojave-desert-tortoise-population-trends>

Berry, K.H., L.J. Allison, A.M. McLuckie, M. Vaughn, and R.W. Murphy. 2021. *Gopherus agassizii*. The IUCN Red List of Threatened Species 2021: e.T97246272A3150871. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T97246272A3150871.en>

[BLM] Bureau of Land Management. 1998a. Las Vegas (Southern Nevada) Field Office Resource Management Plan. Las Vegas, Nevada.

[BLM] U.S. Bureau of Land Management. 1998b. Record of Decision for the Approved Las Vegas Resource Management Plan and Final Environmental Impact Statement. BLM/LV/PL-99/002+1610. Las Vegas Field Office, October 1998.

[CEQ] Council on Environmental Quality. 1997. Considering Cumulative Effects under the National Environmental Policy Act.

- Defenders of Wildlife, Desert Tortoise Preserve Committee, and Desert Tortoise Council. 2020. A Petition to the State of California Fish And Game Commission to move the Mojave desert tortoise from listed as threatened to endangered. Formal petition submitted 11 March 2020. https://defenders.org/sites/default/files/2020-03/Desert%20Tortoise%20Petition%203_20_2020%20Final_0.pdf.
- Gray, M.A., B.G Dickson, K.E. Nussear, T.C. Esque, and T. Chang. 2019. A range-wide model of contemporary, omni-directional connectivity for the threatened Mojave desert tortoise, *Ecosphere* 10(9)e02847. 10.1002/ecs2.2847.
- Jaeger, J. 2005a. Does the configuration of road networks influence the degree to which roads affect wildlife populations? International Conference on Ecology and Transportation 2005 Proceedings, Chapter 5 - Integrating Transportation and Resource Conservation Planning - Landscapes and Road Networks, pages 151-163. August 29, 2005.
- Jaeger, J., J. Bowman, J. Brennan, L. Fahrig, D. Bert, J. Bouchard, N. Charbonneau, K. Frank, B. Gruber, and K. Tluk von Toschanowitz. 2005b. Predicting when animal populations are at risk from roads: an interactive model of road avoidance behavior. *Ecological Modelling* 185 (2005) 329–348.
- Jennings, W.B., and K.H. Berry. 2015. Desert tortoises (*Gopherus agassizii*) are selective herbivores that track the flowering phenology of their preferred food plants. *PLoS ONE* 10(1): e0116716. <https://doi.org/10.1371/journal.pone.0116716>.
- Nussear, K.E., T.C. Esque, R.D. Inman, L. Gass, K.A. Thomas, C.S.A. Wallace, J.B. Blainey, D.M. Miller, and R.H. Webb. 2009. Modeling habitat of the desert tortoise (*Gopherus agassizii*) in the Mojave and parts of the Sonoran Deserts of California, Nevada, Utah, and Arizona. U.S. Geological Survey Open-File Report 2009-1102, 18 p.
- Roedenbeck, I., L. Fahrig, C. Findlay, J. Houlahan, J. Jaeger, N. Klar, S. Kramer-Schadt, and E. van der Grift. 2007. The Rauschholzhausen Agenda for Road Ecology. *Ecology and Society* 12(1): 11. [online]
URL: <http://www.ecologyandsociety.org/vol12/iss1/art11/>
- Tuma, M.W., C. Millington, N. Schumaker, and P. Burnett. 2016. Modeling Agassiz's desert tortoise population response to anthropogenic stressors. *Journal of Wildlife Management* 80(3):414–429; 2016; DOI: 10.1002/jwmg.1044. <https://doi.org/10.1002/jwmg.1044>. <https://wildlife.onlinelibrary.wiley.com/doi/abs/10.1002/jwmg.1044> or <https://www.dropbox.com/s/z27bfcoqyfpz4bd/Tuma%20et%20al%202016%20Modeling%20Agassiz%27s%20desert%20tortoise%20population%20response%20to%20anthropogenic%20stressors.pdf?dl=0>
- [USFWS] U.S. Fish and Wildlife Service. 2014a. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2012 Annual Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. https://www.fws.gov/sites/default/files/documents/2012_Rangewide_Mojave_Desert_Tortoise_Monitoring.pdf

- [USFWS] U.S. Fish and Wildlife Service. 2014b. Status of the desert tortoise and critical habitat. Unpublished report available on the Desert Tortoise Recovery Office's website: "02/10/2014 Status of the Desert Tortoise and Critical Habitat (.704MB PDF)." Reno, NV.
- [USFWS] U.S. Fish and Wildlife Service. 2015. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2013 and 2014 Annual Reports. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2015%20report.%20Rangewide%20monitoring%20report%202013-14.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2016. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2015 and 2016 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2016%20report.%20Rangewide%20monitoring%20report%202015-16.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2017. Status of the desert tortoise and critical habitat (dated 11 October 2017). Unpublished report prepared by the Desert Tortoise Recovery Office of the USFWS. Reno, NV. 24 pages.
- [USFWS] U.S. Fish and Wildlife Service. 2018. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2017 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2018%20report.%20Rangewide%20monitoring%20report%202017.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2019. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2018 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2019%20report.%20Rangewide%20monitoring%20report%202018.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2020. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2019 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. 42 pages. https://www.fws.gov/sites/default/files/documents/2019_Rangewide%20Mojave%20Desert%20Tortoise%20Monitoring.pdf
- [USFWS] U.S. Fish and Wildlife Service. 2022a. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2020 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2022%20report.%20Rangewide%20monitoring%20report%202020.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2022b. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2021 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2022%20report.%20Rangewide%20monitoring%20report%202021.pdf>

Appendix A. Demographic Status and Trend of the Mojave Desert Tortoise (*Gopherus agassizii*)

We provide the following information on the status and trend of the listed population of the desert tortoise to assist the BLM with its analysis of the direct, indirect, and cumulative impacts of the Proposed Project on the Mojave desert tortoise.

BLM's implementation of a conservation strategy for the Mojave desert tortoise in its resource management plans through 2020 has resulted in the following changes in the status for the tortoise throughout its range and in Nevada from 2004 to 2014 (Table 1; USFWS 2015) and 2004 to 2020 (Table 2). There are 17 populations of Mojave desert tortoise described below that occur in the Critical Habitat Units (CHUs) and Tortoise Conservation Areas (TCAs); 14 are on lands managed by the BLM.

The Desert Tortoise Council (Council) has serious concerns about direct, indirect, and cumulative sources of human mortality for the Mojave desert tortoise given the status and trend of the species range-wide, within each of the five recovery units, and within the TCAs that comprise each recovery unit.

Densities of Adult Mojave Desert Tortoises: A few years after listing the Mojave desert tortoise under the Federal Endangered Species Act (FESA), the U.S. Fish and Wildlife Service (USFWS) published a Recovery Plan for the Mojave desert tortoise (USFWS 1994a). It contained a detailed population viability analysis. In this analysis, the minimum viable density of a Mojave desert tortoise population is 10 adult tortoises per mile² (3.9 adult tortoises per km²). This assumed a male-female ratio of 1:1 (USFWS 1994a, page C25) and certain areas of habitat with most of these areas geographically linked by adjacent borders or corridors of suitable tortoise habitat. Populations of Mojave desert tortoises with densities below this density are in danger of extinction (USFWS 1994a, page 32). The revised recovery plan (USFWS 2011) designated five recovery units for the Mojave desert tortoise that are intended to conserve the genetic, behavioral, and morphological diversity necessary for the recovery of the entire listed species (Allison and McLuckie 2018).

Range-wide, densities of adult Mojave desert tortoises declined more than 32% between 2004 and 2014 (Table 1) (USFWS 2015). At the recovery unit level, between 2004 and 2014, densities of adult desert tortoises declined, on average, in every recovery unit except the Northeastern Mojave (Table 1). Adult densities in the Northeastern Mojave Recovery Unit increased 3.1% per year (SE = 4.3%), while the other four recovery units declined at different annual rates: Colorado Desert (-4.5%, SE = 2.8%), Upper Virgin River (-3.2%, SE = 2.0%), Eastern Mojave (-11.2%, SE = 5.0%), and Western Mojave (-7.1%, SE = 3.3%) (Allison and McLuckie 2018). However, the small area and low starting density of the tortoises in the Northeastern Mojave Recovery Unit (lowest density of all Recovery Units) resulted in a small overall increase in the number of adult tortoises by 2014 (Allison and McLuckie 2018). In contrast, the much larger areas of the Eastern Mojave, Western Mojave, and Colorado Desert recovery units, plus the higher estimated initial densities in these areas, explained much of the estimated total loss of adult tortoises since 2004 (Allison and McLuckie 2018).

At the population level, represented by tortoises in the TCAs, densities of 10 of 17 monitored populations of the Mojave desert tortoise declined from 26% to 64% and 11 have densities less than 3.9 adult tortoises per km² (USFWS 2015).

Population Data on Mojave Desert Tortoise: The Mojave desert tortoise was listed as threatened under the FESA in 1990. The listing was warranted because of ongoing population declines throughout the range of the tortoise from multiple human-caused activities. Since the listing, the status of the species has changed. Population numbers (abundance) and densities continue to decline substantially (please see Tables 1 and 2).

Table 1. Summary of 10-year trend data for 5 Recovery Units and 17 CHUs/TCAs for the Mojave desert tortoise, *Gopherus agassizii* (=Agassiz’s desert tortoise). The table includes the area of each Recovery Unit and CHU/TCA, percent of total habitat for each Recovery Unit and CHU/TCA, density (number of breeding adults/km² and standard errors = SE), and the percent change in population density between 2004-2014. Populations below the viable level of 3.9 adults/km² (10 adults per mi²) (assumes a 1:1 sex ratio) and showing a decline from 2004 to 2014 are in red (Allison and McLuckie 2018, USFWS 2015).

Recovery Unit Designated CHU/TCA	Surveyed area (km ²)	% of total habitat area in Recovery Unit & CHU/TCA	2014 density/km ² (SE)	% 10-year change (2004– 2014)
Western Mojave, CA	6,294	24.51	2.8 (1.0)	-50.7 decline
Fremont-Kramer	2,347	9.14	2.6 (1.0)	-50.6 decline
Ord-Rodman	852	3.32	3.6 (1.4)	-56.5 decline
Superior-Cronese	3,094	12.05	2.4 (0.9)	-61.5 decline
Colorado Desert, CA	11,663	45.42	4.0 (1.4)	-36.25 decline
Chocolate Mtn AGR, CA	713	2.78	7.2 (2.8)	-29.77 decline
Chuckwalla, CA	2,818	10.97	3.3 (1.3)	-37.43 decline
Chemehuevi, CA	3,763	14.65	2.8 (1.1)	-64.70 decline
Fenner, CA	1,782	6.94	4.8 (1.9)	-52.86 decline
Joshua Tree, CA	1,152	4.49	3.7 (1.5)	+178.62 increase
Pinto Mtn, CA	508	1.98	2.4 (1.0)	-60.30 decline
Piute Valley, NV	927	3.61	5.3 (2.1)	+162.36 increase
Northeastern Mojave	4,160	16.2	4.5 (1.9)	+325.62 increase
Beaver Dam Slope, NV, UT, AZ	750	2.92	6.2 (2.4)	+370.33 increase
Coyote Spring, NV	960	3.74	4.0 (1.6)	+ 265.06 increase
Gold Butte, NV & AZ	1,607	6.26	2.7 (1.0)	+ 384.37 increase
Mormon Mesa, NV	844	3.29	6.4 (2.5)	+ 217.80 increase
Eastern Mojave, NV & CA	3,446	13.42	1.9 (0.7)	-67.26 decline
El Dorado Valley, NV	999	3.89	1.5 (0.6)	-61.14 decline
Ivanpah Valley, CA	2,447	9.53	2.3 (0.9)	-56.05 decline
Upper Virgin River	115	0.45	15.3 (6.0)	-26.57 decline
Red Cliffs Desert	115	0.45	15.3 (6.0)	-26.57 decline
Total amount of land	25,678	100.00		-32.18 decline

Density of Juvenile Mojave Desert Tortoises: Survey results indicate that the proportion of juvenile desert tortoises has been decreasing in all five recovery units since 2007 (Allison and McLuckie 2018). The probability of encountering a juvenile tortoise was consistently lowest in the Western Mojave Recovery Unit. Allison and McLuckie (2018) provided reasons for the decline in juvenile desert tortoises in all recovery units. These included decreased food availability for adult female tortoises resulting in reduced clutch size, decreased food availability resulting in increased mortality of juvenile tortoises, prey switching by coyotes from mammals to tortoises, and increased abundance of common ravens that typically prey on smaller desert tortoises.

Declining adult tortoise densities through 2014 have left the Eastern Mojave adult numbers at 33% (a 67% decline of their 2004 levels) (Allison and McLuckie 2018, USFWS 2015). Such steep declines in the density of adults are only sustainable if there are suitably large improvements in reproduction and juvenile growth and survival. However, the proportion of juveniles has not increased anywhere in the range of the Mojave desert tortoise since 2007, and in the Eastern Mojave Recovery Unit the proportion of juveniles in 2014 declined from 14 to 11 percent (a 21% decline) of their representation since 2007 (Allison and McLuckie 2018).

The USFWS and Utah Division of Wildlife Resources have continued to collect density data on the Mojave desert tortoise since 2014. The results are provided in Table 2 along with the analysis USFWS (2015) conducted for tortoise density data from 2004 through 2014. These data show that adult tortoise densities in most Recovery Units continued to decline in density since the data collection methodology was initiated in 2004. In addition, in the Northeastern Mojave Recovery Unit that had shown an overall increase in tortoise density between 2004 and 2014, subsequent data indicate a decline in density since 2014 (USFWS 2016, 2018, 2019, 2020, 2022a, 2022b).

Table 2. Summary of data for Agassiz’s desert tortoise, *Gopherus agassizii* (=Mojave desert tortoise) from 2004 to 2021 for the 5 Recovery Units and 17 CHUs/TCAs. The table includes the area of each Recovery Unit and CHU/TCA, percent of total habitat for each Recovery Unit and CHU/TCA, density (number of breeding adults/km² and standard errors = SE), and percent change in population density between 2004-2014 (USFWS 2015). Populations below the viable level of 3.9 breeding individuals/km² (10 breeding individuals per mi²) (assumes a 1:1 sex ratio) (USFWS 1994a, 2015) or showing a decline from 2004 to 2014 are in **red**.

Recovery Unit: Designated CHU/TCA &	% of total habitat area in Recovery Unit & CHU/TCA	2014 density/ km ² (SE)	% 10-year change (2004–2014)	2015 density/ km ²	2016 density/ km ²	2017 density/ km ²	2018 density/ km ²	2019 density/ km ²	2020 density/ km ²	2021 density/ km ²
Western Mojave, CA	24.51	2.8 (1.0)	-50.7 decline							
Fremont-Kramer	9.14	2.6 (1.0)	-50.6 decline	4.5	No data	4.1	No data	2.7	1.7	No data
Ord-Rodman	3.32	3.6 (1.4)	-56.5 decline	No data	No data	3.9	2.5/3.4*	2.1/2.5*	No data	1.9/2.5*
Superior-Cronese	12.05	2.4 (0.9)	-61.5 decline	2.6	3.6	1.7	No data	1.9	No data	No data
Colorado Desert, CA	45.42	4.0 (1.4)	-36.25 decline							
Chocolate Mtn AGR, CA	2.78	7.2 (2.8)	-29.77 decline	10.3	8.5	9.4	7.6	7.0	7.1	3.9
Chuckwalla, CA	10.97	3.3 (1.3)	-37.43 decline	No data	No data	4.3	No data	1.8	4.6	2.6
Chemehuevi, CA	14.65	2.8 (1.1)	-64.70 decline	No data	1.7	No data	2.9	No data	4.0	No data
Fenner, CA	6.94	4.8 (1.9)	-52.86 decline	No data	5.5	No data	6.0	2.8	No data	5.3
Joshua Tree, CA	4.49	3.7 (1.5)	+178.62 increase	No data	2.6	3.6	No data	3.1	3.9	No data

Recovery Unit: Designated CHU/TCA	% of total habitat area in Recovery Unit & CHU/TCA	2014 density/km ² (SE)	% 10- year change (2004– 2014)	2015	2016	2017	2018	2019	2020	2021
Pinto Mtn, CA	1.98	2.4 (1.0)	-60.30 decline	No data	2.1	2.3	No data	1.7	2.9	No data
Piute Valley, NV	3.61	5.3 (2.1)	+162.36 increase	No data	4.0	5.9	No data	No data	No data	3.9
Northeastern Mojave AZ, NV, & UT	16.2	4.5 (1.9)	+325.62 increase							
Beaver Dam Slope, NV, UT, & AZ	2.92	6.2 (2.4)	+370.33 increase	No data	5.6	1.3	5.1	2.0	No data	No data
Coyote Spring, NV	3.74	4.0 (1.6)	+ 265.06 increase	No data	4.2	No data	No data	3.2	No data	No data
Gold Butte, NV & AZ	6.26	2.7 (1.0)	+ 384.37 increase	No data	No data	1.9	2.3	No data	No data	2.4
Mormon Mesa, NV	3.29	6.4 (2.5)	+ 217.80 increase	No data	2.1	No data	3.6	No data	5.2	5.2
Eastern Mojave, NV & CA	13.42	1.9 (0.7)	-67.26 decline							
El Dorado Valley, NV	3.89	1.5 (0.6)	-61.14 decline	No data	2.7	5.6	No data	2.3	No data	No data
Ivanpah Valley, CA	9.53	2.3 (0.9)	-56.05 decline	1.9	No data	No data	3.7	2.6	No data	1.8

Recovery Unit: Designated CHU/TCA	% of total habitat area in Recovery Unit & CHU/TCA	2004 density/ km ²	2014 density/km ² (SE)	% 10-year change (2004–2014)	2015	2016	2017	2018	2019	2020	2021
Upper Virgin River, UT & AZ	0.45		15.3 (6.0)	-26.57 decline							
Red Cliffs Desert**	0.45	29.1 (21.4-39.6)**	15.3 (6.0)	-26.57 decline	15.0	No data	19.1	No data	17.2	No data	
Range-wide Area of CHUs - TCAs/Range-wide Change in Population Status	100.00			-32.18 decline							

*This density includes the adult tortoises translocated from the expansion of the MCAGCC, that is resident adult tortoises and translocated adult tortoises.

**Methodology for collecting density data initiated in 1999.

Abundance of Mojave Desert Tortoises: Allison and McLuckie (2018) noted that because the area available to tortoises (i.e., tortoise habitat and linkage areas between habitats) is decreasing, trends in tortoise density no longer capture the magnitude of decreases in abundance. Hence, they reported on the change in abundance or numbers of the Mojave desert tortoise in each recovery unit (Table 2). They noted that these estimates in abundance are likely higher than actual numbers of tortoises, and the changes in abundance (i.e., decrease in numbers) are likely lower than actual numbers because of their habitat calculation method. They used area estimates that removed only impervious surfaces created by development as cities in the desert expanded. They did not consider degradation and loss of habitat from other sources, such as the recent expansion of military operations (753.4 km² so far on Fort Irwin and the Marine Corps Air Ground Combat Center), intense or large scale fires (e.g., 576.2 km² of critical habitat that burned in 2005), development of utility-scale solar facilities (as of 2015, 194 km² have been permitted) (USFWS 2016), or other sources of degradation or loss of habitat (e.g., recreation, mining, grazing, infrastructure, etc.). Thus, the declines in abundance of Mojave desert tortoise are likely greater than those reported in Table 3.

Table 3. Estimated change in abundance of adult Mojave desert tortoises in each recovery unit between 2004 and 2014 (Allison and McLuckie 2018). Decreases in abundance are in red.

Recovery Unit	Modeled Habitat (km ²)	2004 Abundance	2014 Abundance	Change in Abundance	Percent Change in Abundance
Western Mojave	23,139	131,540	64,871	-66,668	-51%
Colorado Desert	18,024	103,675	66,097	-37,578	-36%
Northeastern Mojave	10,664	12,610	46,701	34,091	270%
Eastern Mojave	16,061	75,342	24,664	-50,679	-67%
Upper Virgin River	613	13,226	10,010	-3,216	-24%
Total	68,501	336,393	212,343	-124,050	-37%

Habitat Availability: Data on population density or abundance does not indicate population viability. The area of protected habitat or reserves for the subject species is a crucial part of the viability analysis along with data on density, abundance, and other population parameters. In the Desert Tortoise (Mojave Population) Recovery Plan (USFWS 1994a), the analysis of population viability included population density and size of reserves (i.e., areas managed for the desert tortoise) and population numbers (abundance) and size of reserves. The USFWS Recovery Plan reported that as population densities for the Mojave desert tortoise decline, reserve sizes must increase, and as population numbers (abundance) for the Mojave desert tortoise decline, reserve sizes must increase (USFWS 1994a). In 1994, reserve design (USFWS 1994a) and designation of critical habitat (USFWS 1994b) were based on the population viability analysis from numbers (abundance) and densities of populations of the Mojave desert tortoise in the early 1990s. Inherent in this analysis is that the lands be managed with reserve level protection (USFWS 1994a, page 36) or ecosystem protection as described in section 2(b) of the FESA, and that sources of mortality be reduced so recruitment exceeds mortality (that is, $\lambda > 1$)(USFWS 1994a, page C46).


Habitat loss would also disrupt the prevailing population structure of this widely distributed species with geographically limited dispersal (isolation by resistance Dutcher et al. 2020). Allison and McLuckie (2018) anticipate an additional impact of this habitat loss/degradation is decreasing

resilience of local tortoise populations by reducing demographic connections to neighboring populations (Fahrig 2007). Military and commercial operations and infrastructure projects that reduce tortoise habitat in the desert are anticipated to continue (Allison and McLuckie 2018) as are other sources of habitat loss/degradation.

Allison and McLuckie (2018) reported that the life history of the Mojave desert tortoise puts it at greater risk from even slightly elevated adult mortality (Congdon et al. 1993; Doak et al. 1994), and recovery from population declines will require more than enhancing adult survivorship (Spencer et al. 2017). The negative population trends in most of the TCAs for the Mojave desert tortoise indicate that this species is on the path to extinction under current conditions (Allison and McLuckie 2018). They state that their results are a call to action to remove ongoing threats to tortoises from TCAs, and possibly to contemplate the role of human activities outside TCAs and their impact on tortoise populations inside them.

Densities, numbers, and habitat for the Mojave desert tortoise declined between 2004 and 2014 and densities continue to decline in most Recovery Units since 2014. As reported in the population viability analysis, to improve the status of the Mojave desert tortoise, reserves (area of protected habitat) must be established and managed. When densities of tortoises decline, the area of protected habitat must increase. When the abundance of tortoises declines, the area of protected habitat must increase. We note that the Desert Tortoise (Mojave Population) Recovery Plan was released in 1994 and its report on population viability and reserve design was reiterated in the 2011 Revised Recovery Plan as needing to be updated with current population data (USFWS 2011, p. 83). With lower population densities and abundance, a revised population viability analysis would show the need for greater areas of habitat to receive reserve level of management for the Mojave desert tortoise. In addition, we note that none of the recovery actions that are fundamental tenets of conservation biology has been implemented throughout most or all of the range of the Mojave desert tortoise.

IUCN Species Survival Commission: The Mojave desert tortoise is now on the list of the world’s most endangered tortoises and freshwater turtles. It is in the top 50 species. The International Union for Conservation of Nature’s (IUCN) Species Survival Commission, Tortoise and Freshwater Turtle Specialist Group, now considers Mojave desert tortoise to be Critically Endangered (Berry et al. 2021). As such, it is a “species that possess an extremely high risk of extinction as a result of rapid population declines of 80 to more than 90 percent over the previous 10 years (or three generations), a current population size of fewer than 50 individuals, or other factors.” It is one of three turtle and tortoise species in the United States to be critically endangered. This designation is more grave than endangered.



Literature Cited in Appendix A

- Allison, L.J. and A.M. McLuckie. 2018. Population trends in Mojave desert tortoises (*Gopherus agassizii*). *Herpetological Conservation and Biology* 13(2):433–452.
- Berry, K.H., L.J. Allison, A.M. McLuckie, M. Vaughn, and R.W. Murphy. 2021. *Gopherus agassizii*. The IUCN Red List of Threatened Species 2021: e.T97246272A3150871. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T97246272A3150871.en>
- Congdon, J.D., A.E. Dunham, and R.C. van Loeben Sels. 1993. Delayed sexual maturity and demographics of Blanding's Turtles (*Emydoidea blandingii*): implications for conservation and management of long-lived organisms. *Conservation Biology* 7:826–833.
- Doak, D., P. Karieva, and B. Klepetka. 1994. Modeling population viability for the Desert Tortoise in the Western Mojave. *Ecological Applications* 4:446–460.
- Dutcher, K.E., A.G. Vandergast, T.C Esque, A. Mittelberg, M.D. Matocq, J.S. Heaton, and K.E. Nussear. 2020. Genes in space: what Mojave desert tortoise genetics can tell us about landscape connectivity. *Conservation Genetics* 21:289–303(2020).
- Fahrig, L. 2007. Non-optimal animal movement in human-altered landscapes. *Functional Ecology* 21:1003–1015.
- Murphy, R.W., K.H. Berry, T. Edwards, and A.M. McLuckie. 2007. A genetic assessment of the recovery units for the Mojave population of the Desert Tortoise, *Gopherus agassizii*. *Chelonian Conservation and Biology* 6:229–251.
- Murphy, R.W., K.H. Berry, T. Edwards, A.E. Leviton, A. Lathrop, and J. D. Riedle. 2011. The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation. *ZooKeys* 113: 39–71. doi: 10.3897/zookeys.113.1353.
- Spencer, R.-J., J.U. Van Dyke, and M.B. Thompson. 2017. Critically evaluating best management practices for preventing freshwater turtle extinctions. *Conservation Biology* 31:1340–1349.
- Turtle Conservation Coalition. 2018. Turtles in Trouble: The World's 25+ Most Endangered Tortoises and Freshwater Turtles. www.iucn-tftsg.org/trouble.
- [USFWS] U.S. Fish and Wildlife Service. 1994a. Desert tortoise (Mojave population) Recovery Plan. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon. 73 pages plus appendices.

- [USFWS] U.S. Fish and Wildlife Service. 1994b. Endangered and threatened wildlife and plants; determination of critical habitat for the Mojave population of the desert tortoise. Federal Register 55(26):5820-5866. Washington, D.C.
- [USFWS] U.S. Fish and Wildlife Service. 2011. Revised Recovery Plan for the Mojave Population of the Desert Tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service, California and Nevada Region, Sacramento, California.
- [USFWS] U.S. Fish and Wildlife Service. 2015. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2013 and 2014 Annual Reports. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2015%20report.%20Rangewide%20monitoring%20report%202013-14.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2016. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2015 and 2016 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2016%20report.%20Rangewide%20monitoring%20report%202015-16.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2018. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2017 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2018%20report.%20Rangewide%20monitoring%20report%202017.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2019. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2018 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2019%20report.%20Rangewide%20monitoring%20report%202018.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2020. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2019 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. 42 pages. https://www.fws.gov/sites/default/files/documents/2019_Rangewide%20Mojave%20Desert%20Tortoise%20Monitoring.pdf
- [USFWS] U.S. Fish and Wildlife Service. 2022a. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2020 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2022%20report.%20Rangewide%20monitoring%20report%202020.pdf>
- [USFWS] U.S. Fish and Wildlife Service. 2022b. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2021 Annual Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. <https://www.fws.gov/sites/default/files/documents/USFWS.2022%20report.%20Rangewide%20monitoring%20report%202021.pdf>

Appendix B. Bibliography on road impacts in desert ecosystems

- Aber, J.D., K.J. Nadelhoffer, P. Steudler, and J.M. Melillo. 1989. Nitrogen Saturation in Northern Forest Ecosystems. *BioScience* 39(6):8-386
- Allen, E.B., Rao, L.E., Steers, R.J., Bytnerowicz, A., and Fenn, M.E., 2009, Impacts of atmospheric nitrogen deposition on vegetation and soils at Joshua Tree National Pages, in Webb, R.H., Fenstermaker, L.F., Heaton, J.S., Hughson, D.L., McDonald, E.V., and Miller, D.M. (eds.), *The Mojave Desert: ecosystem processes and sustainability*: Reno, University of Nevada Press, .p. 78–100.
- Arnold, R. 2011. Focused desert tortoise survey, Lucerne Valley Desert View Ranch generating facility. APN 0435-083-39 & -435-132-01, San Bernardino County. RCA Associated, Hesperia, CA.
- Avery, H.W. 1997. Effects of cattle grazing on the desert tortoise, *Gopherus agassizii*: Nutritional and behavioral interactions. Pages 13-20 in J. Van Abbema (ed.), *Proceedings of the International Conference on Conservation, Restoration, and Management of Tortoises and Turtles*. New York Turtle and Tortoise Society, New York.
- Avery, H.W. 1998. Nutritional ecology of the desert tortoise (*Gopherus agassizii*.) in relation to cattle grazing in the Mojave Desert. Ph.D. dissertation, University of California, Los Angeles.
- Beacon Solar. 2008. Application for Incidental Take of Threatened and Endangered Species. Application to California Department of Fish and Game by Beacon Solar, LLC, 700 Universe Boulevard, Juno Beach, FL.
- Beazley, K.F., T.V. Snaith, F. Mackinnin, and D. Colville. 2004. Road density and potential impacts on wildlife species such as American moose in mainland Nova Scotia. *Proc. N.S. Inst. Sci.* (2004) Volume 42, Part 2, pp. 339-357.
- Belnap, J. 1996. Soil surface disturbances in cold deserts: effects on nitrogenase activity in cyanobacterial-lichen soil crusts. *Biol Fertil Soils* (1996) 23:362-367.
- Berry, K.H. 1974. Desert tortoise relocation project: Status report for 1972. California Department of Transportation
- Berry, K.H. 1990. The status of the desert tortoise in California in 1989. Draft report. U.S. Bureau of Land Management, Riverside, California.
- Berry, K.H., and L.L. Nicholson. 1984b. A summary of human activities and their impacts on desert tortoise populations and habitat in California. Chapter 3 in K.H. Berry (ed.), *The status of the desert tortoise (*Gopherus agassizii*) in the United States*. Desert Tortoise Council Report to the U.S. Fish and Wildlife Service. Order No. 11310-0083-81.

- Berry, K.H., F.G. Hoover, and M. Walker. 1996. The effects of poaching desert tortoises in the western Mojave Desert; evaluation of landscape and local impacts. Proceedings of the Desert Tortoise Council Symposium 1996:45.
- Berry, K.H., K. Keith, and T. Bailey. 2008. Status of the desert tortoise in Red Rock Canyon State Park. California Fish and Game 94(2):98-118.
- Berry, K.H., J. L. Yee, A.A. Coble, W.M. Perry, and T.A. Shields. 2013. Multiple factors affect a population of Agassiz's desert tortoise (*Gopherus agassizii*) in the northwestern Mojave Desert. Herpetological Monographs, 27, 2013, 87–109.
- Berry, K.H., L.M. Lyren, J.L. Yee, and T.Y. Bailey. 2014. Protection benefits desert tortoise (*Gopherus agassizii*) abundance: the influence of three management strategies on a threatened species. Herpetological Monographs, 28 2014, 66–92.
- Berry, K.H., L.M. Lyren, J.S. Mack, L.A Brand, and D.A.Wood. 2016. Desert tortoise annotated bibliography, 1991–2015: U.S. Geological Survey Open-File Report 2016-1023, 312 p., <http://dx.doi.org/10.3133/ofr20161023>.
- Boarman, W.I. 2002. Threats to desert tortoise populations: a critical review of the literature. Unpublished Report, prepared for the West Mojave Planning Team and the Bureau of Land Management. 86 pp.
- Boarman, W.I., and K.H Berry. 1995. Common ravens in the southwestern United States, 1968-92. In: Our Living Resources: A Report to the Nation on the Distribution, Abundance, and Health of U.S. Plants, Animals, and Ecosystems. Edward T. LaRoe, Gaye S. Farris, Catherine E. Puckett, Peter D. Doran, and Michael J. Mac, editors. U.S. Department of the Interior, National Biological Service.
- Boarman, W.I., R.J. Camp, M. Hagan, W. Deal. 1995. Raven abundance at anthropogenic resources in the western Mojave Desert, California. Report to Edwards Air Force Base, California.
- Boarman, W.I., and M. Sazaki. 1996. Highway mortality in desert tortoises and small vertebrates: success of barrier fences and culverts. Proceedings: Florida Department of Transportation/Federal Highway Administration Transportation-Related Wildlife Mortality Seminar. Evink, G., Ziegler, D., Garrett, P. and Berry, J. (Eds). pp. 169–173.
- Boarman, W.I., and Sazaki, M., 2006, A highway's road-effect zone for desert tortoises (*Gopherus agassizii*): Journal of Arid Environments, v. 65, p. 94–101.
- Boarman, W.I., Sazaki, M., Jennings, B., 1997. The effects of roads, barrier fences and culverts on desert tortoise populations in California, USA. In: Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles — An International Conference, pp. 54–58.

- Boarman, W.I., M.L. Beigel, G.C. Goodlett, and M. Sazaki. 1998. A passive integrated transponder system for tracking animal movements. *Wildlife Society Bulletin* 26, 886-891.
- Boarman, W.I., W.B. Kristan, W.C. Webb, and H.D. Chamblin. 2005. Raven ecology in the Mojave Desert at Edwards Air Force Base: final report. U.S. Geological Survey, Western Ecological Research Center, Sacramento, California.
- Boarman, W.I., and W.B. Kristan. 2006. Evaluation of evidence supporting the effectiveness of desert tortoise recovery actions. Scientific Investigations Report 2006-5143. U.S. Geological Survey, Western Ecological Research Center, Sacramento, CA.
- Boarman, W.I., Patten, M.A., Camp, R.J., and Collis, S.J., 2006, Ecology of a population of subsidized predators: common ravens in the central Mojave Desert, California: *Journal of Arid Environments*, v. 67, p. 248-261.
- Bouchard, J., A. T. Ford, F. Eigenbrod, and L. Fahrig. 2009. Behavioral response of northern leopard frogs (*Rana pipens*) to roads and traffic: implications for population persistence. *Ecology and Society* 14(2): 23. [online] URL: <http://www.ecologyandsociety.org/vol14/iss2/art23/>.
- Bratzel, S., and R. Tellermann. 2005. Mobilität und Verkehr. *Informationen zur politischen Bildung* 287(2):44-51.
- Brocke, R.H., J.P. O'Pezio, K.A. Gustafson. 1988 A forest management scheme mitigating impact of road networks on sensitive wildlife species. In: R.M. Degraaf and W.M. Healy (eds): *Is forest fragmentation a management issue in the northeast? GTR-NE-140*, U.S. Department of Agriculture, Forest Service, Northeastern Forest Experimental Station, Radnor, PA: 13-17.
- Brocke, R.H., J.P. O'Pezio, K.A. Gustafson. 1988 A forest management scheme mitigating impact of road networks on sensitive wildlife species. In: R.M. Degraaf and W.M. Healy (eds): *Is forest fragmentation a management issue in the northeast? GTR-NE-140*, U.S. Department of Agriculture, Forest Service, Northeastern Forest Experimental Station, Radnor, PA: 13-17.
- Brooks, M.L., 1995, Benefits of protective fencing to plant and rodent communities of the western Mojave Desert, California: *Environmental Management*, v. 19, p. 65-74.
- Brooks, M.L., 1999, Alien annual grasses and fire in the Mojave Desert: *Madroño*, v. 46, p. 13-19.
- Brooks, M.L., 2003, Effects of increased soil nitrogen on the dominance of alien annual plants in the Mojave Desert: *Journal of Applied Ecology*, v. 40, p. 344-353.
- Brooks, M.L., T.C. Esque, and J.R. Matchett. 2003. Current status and management of alien plants and fire in desert tortoise habitat. *Desert Tortoise Council Symposium*, February 21-23, 2003.

- Brooks, M.L. 2009, Spatial and temporal distribution of non-native plants in upland areas of the Mojave Desert, in Webb, R.H., Fenstermaker, L.F., Heaton, J.S., Hughson, D.L., McDonald, E.V., and Miller, D.M., eds., *The Mojave Desert—Ecosystem processes and sustainability*: Reno, University of Nevada Press, p. 101–124.
- Brooks, M.L. and K.H. Berry. 1999. Ecology and management of alien annual plants in the California deserts. *Calif. Exotic Pest Plant Newsl.* 7(3/4):4-6.
- Brooks, M.L., and Berry, K.H., 2006, Dominance and environmental correlates of alien annual plants in the Mojave Desert, USA: *Journal of Arid Environments*, v. 67, p. 100–124.
- Brooks, M.L., and Esque, T.C., 2002, Alien plants and fire in desert tortoise (*Gopherus agassizii*) habitat of the Mojave and Colorado Deserts: *Chelonian Conservation and Biology*, v. 4, p. 330–340.
- Brooks, M.L., C.M. D’Antonio, D.M. Richardson, J. B. Grace, J.E. Kelley, J. M. Ditomaso, R.J. Hobbs, M. Pellant, And D. Pyke. 2004. Effects of Invasive Alien Plants on Fire Regimes. *Bioscience/ Vol. 54 No. 7: 677-688.* July 2004.
- Brooks, M.L. and B. Lair. 2005. Ecological Effects of Vehicular Routes in a Desert Ecosystem. Report prepared for the United States Geological Survey, Recoverability and Vulnerability of Desert Ecosystems Program (<http://geography.wr.usgs.gov/mojave/rvde>).
- Brooks, M.L. and B. M. Lair.2009. Ecological effects of vehicular routes in a desert ecosystem. In: R.H. Webb, L.F. Fenstermaker, J.S. Heaton, D.L. Hughson, E.V. McDonald, and D.M. Miller (eds.). *The Mojave Desert: Ecosystem Processes and Sustainability*. University of Arizona Press. Tucson, AZ.
- Brooks, M.L., and Matchett, J.R. 2006. Spatial and temporal patterns of wildfires in the Mojave Desert, 1980–2004. *Journal of Arid Environments* Volume 67, Supplement, 2006, Pages 148-164.
- Brown, D.E., and R.A. Minnich. 1986. Fire and changes in creosote bush scrub of the western Sonoran desert, California. *American Naturalist* 116(2):411-422.
- Bureau of Land Management. U. S. Fish and Wildlife Service, and California Department of Fish and Game. 1989. Environmental assessment for selected control of the common raven to reduce desert tortoise predation in the Mojave Desert, California. Bureau of Land Management, U. S. Fish and Wildlife Service, and California Department of Fish and Game.
- Bureau of Land Management. 1993. Final Rand Mountains–Fremont Valley Management Plan. A Sikes Act Plan. Bureau of Land Management, Ridgecrest Resource Area, California.

- Bureau of Land Management. 1998. The California Desert Conservation Area Plan 1980, as amended. U.S. Department of the Interior, Bureau of Land Management, California.
- Bureau of Land Management. 1999. Chapter Two - Desert Tortoise (*Gopherus agassizii*). Working draft for West Mojave Plan. September 22, 1999. https://www.blm.gov/ca/pdfs/cdd_pdfs/Ch2_9-22-99.pdf
- Bureau of Land Management. 2002. Proposed Northern and Eastern Mojave Desert management plan - Final Environmental Impact Statement – Volumes 1 and 2. California Desert District, Riverside, CA.
- Bureau of Land Management, County of San Bernardino, and City of Barstow. 2005. Proposed West Mojave Plan Final Environmental Impact Report and Statement. BLM/CA/ES-2004-005 + 1790 -1600. Moreno Valley, CA.
- Bureau of Land Management. 2006. Record of decision for the West Mojave Plan. California Desert District, Moreno Valley, CA.
- Bureau of Land Management. 2018a. West Mojave Route Network Project Draft California Desert Conservation Plan Amendment and Supplemental Environmental Impact Statement for the California Desert District. BLM/CA/DOI-BLM-CA-D080-2018-0008-EIS. January 2018. Moreno Valley, CA.
- Bureau of Land Management. 2018b. Stoddard Valley OHV Area. BLM website accessed May 30, 2018. <https://www.blm.gov/visit/stoddard-valley-ohv-area>
- Burge, B.L. 1977. Daily and seasonal behavior, and areas utilized by the desert tortoise, *Gopherus agassizii*, in southern Nevada. Proceedings of the Desert Tortoise Council Symposium 1977:59-94.
- Bury, R.B., and Luckenbach, R.A., 2002, Comparison of desert tortoise (*Gopherus agassizii*) populations in an unused and off-road vehicle area in the Mojave Desert: Chelonian Conservation and Biology, v. 4, p. 457–463.
- Caid, N., P. Crist, R. Gilbert, and P. Wiederkehr. 2002. Environmentally sustainable transport: concept, goal and strategy—the OECD’s EST Project. Proceedings of the Institution of Civil Engineers, Transport 153(4):219-226.
- California Turtle and Tortoise Club. 2002. Western Rand Mountains ACEC vehicle closure. <https://tortoise.org/conservation/randacec.html>
- Carr, L. W., and L. Fahrig. 2001. Effect of road traffic on two amphibian species of different vagility. Conservation Biology 15(4):1071-1078.

- Charis Corporation. 2005. Supplemental Final Environmental Impact Statement - Proposed Addition of Maneuver Training Land at Fort Irwin, CA. August 2005. Prepared for the U.S. Army National Training Center, Fort Irwin, California.
- D'Antonio, C.M., and Vitousek, P.M., 1992, Biological invasions by exotic grasses, the grass-fire cycle, and global change: *Annual Review of Ecology and Systematics*, v. 23, p. 63–87.
- DeFalco, L.A., Detling, J.K., Tracy, C.R., and Warren, S.D., 2001, Physiological variation among native and exotic winter annual plants associated with microbiotic crusts in the Mojave Desert: *Plant and Soil*, v. 234, p. 1–14.
- Desert Gazette. 2018. El Paso Mountains. <http://digital-desert.com/el-paso-mountains/> (accessed 2018-5-30)
- Doak, D., P. Kareiva, and B. Klepetka. 1994. Modeling population viability for the desert tortoise in the western Mojave Desert. *Ecological Applications* 4:446–460.
- Edwards T., A.E. Karl, M. Vaughn, P.C. Rosen, C.M. Torres, and R.W. Murphy. 2016. The desert tortoise trichotomy: Mexico hosts a third, new sister-species of tortoise in the *Gopherus morafkai*–*G. agassizii* group. *ZooKeys* 562: 131– 158. doi: 10.3897/Zookeys. 562.6124.
- Esque, T.C. 1992. Diet selection of the desert tortoise in the northeast Mojave Desert – FY 1991 update. *Proceedings of the Desert Tortoise Council Symposium* 1992:64-68.
- Esque, T.C. 1994. Diet and diet selection of the desert tortoise (*Gopherus agassizii*) in the northeastern Mojave Desert. Master's Thesis. Colorado State University, Fort Collins.
- Esque, T.C., Schwalbe, C.R., DeFalco, L.A., Duncan, R.B., and Hughes, T.J., 2003, Effects of desert wildfires on desert tortoise (*Gopherus agassizii*) and other small vertebrates: *Southwestern Naturalist*, v. 48, p. 103–111.
- Estrada, J. 2017. Events. *Tortoise Tracks* 37:2 page 1, Summer 2017.
- Fahrig, L., and T. Rytwinski. 2009. Effects of roads on animal abundance: an empirical review and synthesis. *Ecology and Society* 14(1): 21. [online] URL: <http://www.ecologyandsociety.org/vol14/iss1/art21/>
- Federal Highway Administration and California Department of Transportation. 2017. Olancho/Cartago Four-Lane Project on U.S. Highway 395 in Inyo County from 2.1 miles south of LA Aqueduct Bridge (#48-068R) to 0.2 mile south of Ash Creek Bridge (#48-11). Final Environmental Impact Report/ Environmental Assessment with Finding of No Significant Impact and Section 4(f) Evaluation.
- Forman, R. T. T. 2000. Estimate of the area affected ecologically by the road system in the United States. *Conservation Biology* 14(1):31-35.

- Forman, R. T. T., D. Sperling, J. A. Bissonette., A. P. Clevenger, C. D. Cutshal, V. H. Dale, L. Fahrig, R. France, C. R. Goldman, K. Haenue , J. A. Jones, F. J. Swanson, T. Turrentine, and T. C. Winter. 2002. Road ecology—science and solutions. Island Press, Washington, D.C., USA.
- Forman, R.T.T., D.S. Friedman, D. Fitzhenry, J.D. Martin, A.S. Chen, and L.E. Alexander. 1997. Ecological effects of roads: toward three summary indices and an overview of North America. In: Canter K (ed) Habitat fragmentation and infrastructure. Minister of Transport and Public Works and Water Management, Delft, Netherlands, p 40-54.
- Gelbard, J. L., and J. Belnap. 2003. Roads as conduits for exotic plant invasions in a semiarid landscape. *Conservation Biology* 17:420-432.
- Gibbs, J.P., and W.G. Shriver. 2002. Estimating the effects of road mortality on turtle populations. *Conserv. Biol.* 16, 1647–1652.
- Goodlett, G. O. and G. C. Goodlett. 1993. Studies of unauthorized off-highway vehicle activity in the Rand Mountains and Fremont Valley, Kern County, California. *Proc. 1992 Desert Tort. Counc. Symp.* 1993:163-187.
- Gucinski, H., M. Furniss, R. Ziermer, and M. Brookes. 2001. Forest Service roads: a synthesis of scientific information. Gen Tech Rep PNW-GTR-509.1, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR.
- Henen, B.T. 1992. Desert tortoise diet and dietary deficiencies that may limit egg production at Goffs, California. *Proceedings of the Desert Tortoise Council Symposium 1992:97.*
- Hessing, Mark. Botanist for Fort Irwin. E-mail sent to Connie Rutherford, U.S. Fish and Wildlife Service, Ventura Office, regarding off-road vehicle activity on Coolgardie Mesa. June 3, 2006. Cited in: U.S. Fish and Wildlife Service 2008. Lane Mountain milk-vetch (*Astragalus jaegerianus*) 5-Year Review: Summary and Evaluation. U.S. Fish and Wildlife Service, Ventura Fish and Wildlife Office, Ventura, California. June 2008.
- Jaeger, J.A.G., L. Fahrig, and K.C. Ewald. 2005a. Does the configuration of road networks influence the degree to which roads affect wildlife populations? *International Conference on Ecology and Transportation 2005 Proceedings, Chapter 5 - Integrating Transportation and Resource Conservation Planning - Landscapes and Road Networks*, pages 151-163. August 29, 2005.
- Jaeger, J.A.G., J. Bowman, J. Brennan, L. Fahrig, D. Bert, J. Bouchard, N. Charbonneau, K. Frank, B. Gruber, K. Tluk von Toschanowitz. 2005b. Predicting when animal populations are at risk from roads: an interactive model of road avoidance behavior. *Ecological Modelling* 185 (2005) 329–348.

- Jalkotzy, M.G., P.I. Ross, and M.D. Nasserden. 1997. The effects of linear developments on wildlife: a review of selected scientific literature. Arc Wildlife Services Ltd, prepared for Canadian Association of Petroleum Producers, Calgary, Alberta.
- Jennings, B. 1992. Observations on the feeding habits and behavior of desert tortoises at the Desert Tortoise Natural Area, California. Proceeding of the Desert Tortoise Council Symposium 1992:69-81.
- Jennings, W. B. 1993. Foraging ecology of the desert tortoise (*Gopherus agassizii*) in the western Mojave desert. Master's thesis. Arlington, University of Texas: 101 pp.
- Jennings, W. B. 1997. Habitat use and food preferences of the desert tortoise, *Gopherus agassizii*, in the western Mojave Desert and impacts of off-road vehicles. In J. Van Abbema (ed.), Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles-An International Conference, pp. 42-45. New York Turtle and Tortoise Society, New York.
- Karraker, N.E., and J.P. Gibbs. 2011. Contrasting road effect signals in reproduction of long-versus short-lived amphibians. *Hydrobiologia* 664, 213–218.
- Kemp, P.R., and Brooks, M.L., 1998, Exotic species of California deserts: *Fremontia*, v. 26, p. 30–34.
- Kilgo, J.C., R.F. Labisky, and D.E. Fritzen. 1998. Influences of hunting on the behavior of white-tailed deer: implications for conservation of the Florida panther. *Conservation Biology* 12:1359-1364.
- Knight, R.L., and Kawashima, J.Y., 1993, Responses of raven and red-tailed hawk populations to linear right-of-ways: *Journal of Wildlife Management*, v. 57, p. 266–271.
- Knight, R.L., Camp, R.J., Boarman, W.I., and Knight, H.A.L., 1999, Predatory bird populations in the east Mojave Desert, California: *Great Basin Naturalist*, v. 59, p. 331–338.
- LaBerteaux, D.L. 2006. Mustard removal at the Desert Tortoise Research Natural Area, Kern County, California. Report to the Desert Tortoise Preserve Committee, Inc.
- LaRue, E. 1992. Distribution of desert tortoise sign adjacent to Highway 395, San Bernardino County, California. Proceedings of the 1992 Symposium of the Desert Tortoise Council.
- LaRue, E. 1994. Follow-up monitoring report for Stoddard Valley-to-Johnson Valley Point-to-Point Corridor Run. Unpublished report prepared on behalf of the American Motorcyclists Association for the Barstow Resource Area of the Bureau of Land Management.
- LaRue, E. 2008. Latest information on tortoises and other special-status species in Morongo Basin. Morongo Tortoise Update.7-18-2008. Circle Mountain Biological Consultants, Wrightwood, CA. http://www.yucca-valley.org/pdf/general_plan/mb_tortoise_update_july2008.pdf

- LaRue, E. 2014. Mohave Ground Squirrel Trapping Results for Phacelia Wildflower Sanctuary, Los Angeles County, California. <https://www.wildlife.ca.gov/Conservation/Mammals/Mohave-Ground-Squirrel/TAG/BlogPage/4/Month/4/Year/2018>
- Lei, S. A. 2004. Soil compaction from human trampling, biking, and off-road motor vehicle activity in a blackbrush (*Coleogyne ramosissima*) shrubland. *Western North American Naturalist* 64:125-130.
- Loughran, C.L., J.R. Ennen, and J.E. Lovich. 2011. *Gopherus agassizii* (desert tortoise). Burrow collapse. *Herpetological Review* 42(4), 593.
- Lovich, J.E., and Bainbridge, D., 1999, Anthropogenic degradation of the southern California desert ecosystem and prospects for natural recovery and restoration: *Environmental Management*, v. 24, p. 309–326.
- Lovich, J.E., C.B. Yackulic, J. Freilich M. Agha, M. Austin, K.P. Meyer, T.R. Arundel, J. Hansen, M.S. Vamstad, S.A. Root. 2014. Climatic variation and tortoise survival: Has a desert species met its match? *Biological Conservation* 169 (2014) 214–224.
- McLellan, B.N., and D.M Shackleton, 1988. Grizzly bears and resource extraction industries: effects of roads on behavior, habitat use and demography. *J. Appl. Ecol.* 25, 451–460.
- McLuckie, A.M., M.R.M. Bennion, and R.A. Fridell. 2007. Tortoise mortality within the Red Cliffs Desert Reserve following the 2005 wildfire. Utah Division of Wildlife Resource Publication 07-05.
- Medica, P.A., R.B. Bury, and F.B. Turner. 1975. Growth of the desert tortoise (*Gopherus agassizii*) in Nevada. *Copeia* 1975:639-643.
- Minnich, J.E. 1970. Water and electrolyte balance of the desert iguana, *Dipsosaurus dorsalis*, in its native habitat. *Comparative Biochemistry and Physiology* 35:921-933.
- Minnich, J.E. 1979. Comparison of maintenance electrolyte budgets of free-living desert and gopher tortoises (*Gopherus agassizii* and *G. polyphemus*). *Proceedings of the Desert Tortoise Council Symposium 1979* Pp.166-174.
- Murphy, R.W., Berry, K.H., Edwards, T., Leviton, A.E., Lathrop, A., and Riedle, J.D., 2011, The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation: *ZooKeys*, v. 113, p. 39–71.
- Nafus, M.G., T.D. Tuberville, K. A. Buhlmann, and B.D. Todd. 2013. Relative abundance and demographic structure of Agassiz's desert tortoise (*Gopherus agassizii*) along roads of varying size and traffic volume. *Biological Conservation* 162 (2013) 100–106.
- Nagy, K.A. 1972. Water and electrolyte budgets of a free-living desert lizard, *Sauromalus obesus*. *Journal of Comparative Physiology* 79:93-102.

- Nagy, K.A., and P.A. Medica. 1986. Physiological ecology of desert tortoises. *Herpetologica* 42:73-92.
- Nagy, K.A., Henen, B.T., and Vyas, D.B., 1998, Nutritional quality of native and introduced food plants of wild desert tortoises: *Journal of Herpetology*, v. 32, p. 260–267.
- Noss, R. F. 1993. Wildlife corridors. Pages 43-68 in D. S. Smith and P. C. Hellmund, editors. *Ecology of Greenways*. University of Minneapolis Press, Minneapolis, Minnesota.
- Noss, R. F. 1995. Maintaining ecological integrity in representative reserve networks. World Wildlife Fund, Canada.
- [OECD] Organization for Economic Co-operation and Development. 2002. OECD guidelines towards environmentally sustainable transport. OECD Publications, Paris, France.
- Oftedal, O.T. 2002. The nutritional ecology of the desert tortoise in the Mojave and Sonoran deserts. Pages 194-241 in T.R. Van Devender (ed.), *The Sonoran Desert Tortoise; Natural History, Biology and Conservation*. University of Arizona Press, Tucson, Arizona.
- Oftedal, O.T., L.S. Hillard, and D.J. Morafka. 2002. Selective spring foraging by juvenile desert tortoises (*Gopherus agassizii*) in the Mojave Desert—Evidence of an adaptive nutritional strategy: *Chelonian Conservation and Biology*, v. 4, p. 341–352.
- Oftedal, O.T. and M.E. Allen. 1996. Nutrition as a major facet of reptile conservation. *Zoo Biology* 15:491-497.
- Parendes, L.A., and J.A. Jones. 2000. Role of light availability and dispersal in exotic plant invasion along roads and streams in the H. J. Andrews Experimental Forest, Oregon. *Conservation Biology* 14:64.
- Rytwinski, T., and L. Fahrig. 2011. Reproductive rate and body size predict road impacts on mammal abundance. *Ecol. Appl.* 21, 589–600.
- Rytwinski, T., and L. Fahrig. 2012. Do species life history traits explain population responses to roads? A meta-analysis. *Biol. Conserv.* 147, 87–98.
- Roedenbeck, I.A., L. Fahrig, C. S. Findlay, J. E. Houlahan, J.A.G. Jaeger, N. Klar, S. Kramer-Schadt, and E. A. van der Grift. 2007. The Rauschholzhausen Agenda for Road Ecology. *Ecology and Society* 12(1): 11. [online]. <http://www.ecologyandsociety.org/vol12/iss1/art11/>
- Rudis, V.A. 1995. Regional forest fragmentation effects on bottomland hardwood community types and resource values. *Landsc. Ecol.* 10:291-307.

- Sanson, L. 2016. Marines seek plan to move tortoises from Johnson Valley. Hi-Desert Star September 8, 2016. http://www.hidesertstar.com/news/article_c51696c6-7609-11e6-847d-03224974e42a.html
- Sazaki, M., W.I. Boarman, G. Goodlett, and T. Okamoto. 1995. Risk associated with long-distance movement by desert tortoises. Proceedings of the Desert Tortoise Council 1994 Symposium. pp. 33–48.
- W.H. Schlesinger, and C.S. Jones. 1984. The Comparative Importance of Overland Runoff and Mean Annual Rainfall to Shrub Communities of the Mojave Desert. Botanical Gazette 1984 145(1): 116-124.
- Sharifi, M.R., A.C. Gibson, and P.W. Rundel. 1997. Surface Dust Impacts on Gas Exchange in Mojave Desert Shrubs. Journal of Applied Ecology, 34(4)(Aug., 1997):837-846.
- Sherwood, B., D. Cutler, and J. A. Burton. 2002. Wildlife and roads—the ecological impact. Imperial College Press, London, UK.
- Spellerberg, I. F. 2002. Ecological effects of roads. Land Reconstruction and Management Series, Volume 2. Science Publishers, Enfield, UK.
- Tierra Madre Consultants. 1991. Biological assessment for Lancaster City and Planning Area: Relative density surveys for desert tortoises and cumulative human impact evaluations for Mohave ground squirrel habitat. Report prepared by Ed LaRue for City of Lancaster. Tierra Madre Consultants, Riverside, CA.
- Tracy, C.R., L.C. Zimmerman, C. Tracy, K.D. Bradley, and K. Castle. 2006. Rates of food passage in the digestive tract of young desert tortoises: Effects of body size and diet quality. Chelonian Conservation and Biology: December 2006, Vol. 5, No. 2, pp. 269-273.
- Tratz, W.M. 1978. Postfire vegetational recovery, productivity and herbivore utilization of a chaparral-desert ecotone. Master's Thesis. California State University, Los Angeles.
- Tratz, W.M., and R.J. Vogl. 1977. Postfire vegetational recovery, productivity and herbivore utilization of a chaparral-desert ecotone. Pages 426-430 in H.A. Mooney and C.E. Conrad (eds.), Proceedings of Symposium on Environmental Consequences of Fire and Fuel Management in Mediterranean Ecosystems. USDA Forest Service General Technical Report WO-3.
- Trombulak, S. C., and C. A. Frissell. 2000. Review of ecological effects of roads on terrestrial and aquatic communities. Conservation Biology 14:18–30.
- Turner, F.B., P. Hayden, B.L. Burge, and J.B. Roberson. 1986. Egg production by the desert tortoise (*Gopherus agassizii*) in California. Herpetologica 42:93-104.

- Turner, F.B., K.H. Berry, D.C. Randall, and G.C. White. 1987. Population ecology of the desert tortoise at Goffs, California, 1983-1986. Report to Southern California Edison Co., Rosemead, California.
- Turtle Conservation Coalition. 2018. Turtles in Trouble: The World's 25+ Most Endangered Tortoises and Freshwater Turtles. www.iucn-tftsg.org/trouble
- Umweltbundesamt (UBA). 2003. Reduzierung der Flächeninanspruchnahme durch Siedlung und Verkehr. Materialienband. Umweltbundesamt Texte 90/03, Berlin, Germany. Available online at: <http://www.umweltdaten.de/publikationen/fpdf-l/2587.pdf>.
- Underhill, J. E., and P. G. Angold. 2000. Effects of roads on wildlife in an intensively modified landscape. *Environmental Reviews* 8:21-39.
- U.S. District Court. 2011. Order re: remedy. Case 3:06-cv04884-SI. Center for Biological Diversity, et al., Plaintiffs v. BLM. United States District Court for the Northern District of California, USA. As cited in Berry, K.H., L.M. Lyren, J.L. Yee, and T.Y. Bailey. 2014. Protection benefits desert tortoise (*Gopherus agassizii*) abundance: the influence of three management strategies on a threatened species. *Herpetological Monographs*, 28 2014, 66–92.
- U.S. Ecology. 1989. Proponent's Environmental Assessment. California Low-Level Radioactive Waste Disposal Site. Appendices K and M, Volume II.
- U.S. Fish and Wildlife Service. 1994a. Endangered and threatened wildlife and plants; determination of critical habitat for the Mojave population of the desert tortoise. *Federal Register* 55(26):5820-5866. Washington, D.C.
- U.S. Fish and Wildlife Service. 1994b. Desert tortoise (Mojave population) Recovery Plan. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon. 73 pages plus appendices.
- U.S. Fish and Wildlife Service. 2008. Lane Mountain milk-vetch (*Astragalus jaegerianus*) 5-Year Review: Summary and Evaluation. Ventura Fish and Wildlife Office, Ventura, California.
- U.S. Fish and Wildlife Service. 2011a. Revised recovery plan for the Mojave population of the desert tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California. 222 pp.
- U.S. Fish and Wildlife Service. 2011b. Biological Opinion on Mojave Solar, LLC's Mojave Solar Project, San Bernardino County, California (8-8-11-F-3). Ventura Fish and Wildlife Office, Ventura, CA.
- U.S. Fish and Wildlife Service 2014a. 12-month finding on a petition to reclassify *Astragalus jaegerianus* as a threatened Species. 79 *Federal Register* 25084-25092, Friday, May 2, 2014.

- U.S. Fish and Wildlife Service. 2014b. Determination of threatened status for the western distinct population segment of the yellow-billed cuckoo (*Coccyzus americanus*); Final Rule. 79 Federal Register 59992-60038.
- U.S. Fish and Wildlife Service. 2014c. Designation of critical habitat for the western distinct population segment of the yellow-billed cuckoo; Proposed Rule. 29 Federal Register 48548-48652.
- U.S. Fish and Wildlife Service. 2015. Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2013 and 2014 Annual Reports. Report prepared by Linda Allison for the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- van der Ree R., J. A. G. Jaeger, E. A. van der Grift, and A. P. Clevenger. 2011. Effects of roads and traffic on wildlife populations and landscape function: Road ecology is moving toward larger scales. *Ecology and Society* 16(1): 48. [online] URL: <http://www.ecologyandsociety.org/vol16/iss1/art48/>
- von Seckendorff Hoff, K., and Marlow, R.W. 2002. Impacts of vehicle road traffic on desert tortoise populations with consideration of conservation of tortoise habitat in southern Nevada. *Chelonian Conservation and Biology* 4:449–456.
- Zimmerman, L.C., Espinoza, R.E., and Barber, A.M., 2006a, The importance of physiological ecology in conservation biology: *Integrative and Comparative Biology*, v. 46, p. 1,191–1,205.

Appendix C. Habitat Restoration Bibliography for the Mojave Desert

- Abella, S.A., K.H. Berry, and S. Ferrazzano. 2023. Techniques for Restoring Damaged Mojave and Western Sonoran Habitats, Including Those for Threatened Desert Tortoises and Joshua Trees. *Desert Plants Journal* Volume 38, Number 2, April 2023, pages 4-52.
- Abella, S.R., and A.C. Newton. 2009. A Systematic Review of Species Performance and Treatment Effectiveness for Revegetation in the Mojave Desert, USA. In: *Arid Environments and Wind Erosion*, A. Fernandez-Bernal and M. A. De La Rosa (eds.). Nova Science Publishers. ISBN: 978-1-60692-411-2.
- Abella, S.R., L.P. Chiquoine, E. Cayenne Engel, K.E. Kleinick, and F.S. Edwards. 2015. Enhancing Quality of Desert Tortoise Habitat: Augmenting Native Forage and Cover Plants. *Journal of Fish and Wildlife Management* 6(2): 278-289.
- Abella, S.R., L.P. Chiquoine, A.C. Newton, and C. H. Vanier. 2015. Restoring a desert ecosystem using soil salvage, revegetation, and irrigation. *Journal of Arid Environments* 115 (April 2015): 44-52.
- Abella, S.R., and K.H. Berry. 2016. Enhancing and restoring habitat for the desert tortoise. *Journal of Fish and Wildlife Management* 7(1):255-279; e1944-687X. doi: 10.3996/052015-JFWM-046.
- Bowker, M.A., M. E. Miller, J. Belnap, T.D. Sisk, and N.C. Johnson. 2008. Prioritizing Conservation Effort through the Use of Biological Soil Crusts as Ecosystem Function Indicators in an Arid Region. *Conservation Biology* 22(6):1533-1543.
- Chiquoine, L.P., S.R. Abella, and M.A. Bowker. 2016. Rapidly restoring biological soil crusts and ecosystem functions in a severely disturbed desert ecosystem. *Ecological Applications* 26(4): 1260-1272. <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/15-0973>
- Esque, T.C., L. A. DeFalco, G.L. Tyree, K. K. Drake, K.E. Nussear, and J.S. Wilson. 2021. Priority Species Lists to Restore Desert Tortoise and Pollinator Habitats in Mojave Desert Shrublands. *Natural Areas Journal* 41(2):145-158.
- Jennings, W.B., and K.H. Berry. 2015. Desert tortoises (*Gopherus agassizii*) are selective herbivores that track the flowering phenology of their preferred food plants. *PLoS ONE* 10(1): e0116716. <https://doi.org/10.1371/journal.pone.0116716>.
- Rader, A.J. 2019. Assessing Minimal-Input Restoration Strategies for Desert Soil and Vegetation Restoration. Master of Science Thesis, School of Life Sciences, University of Nevada, Las Vegas.
- Shryock, D.F., L.A. DeFalco, and T.C. Esque. 2022. Seed Menus: An integrated decision-support framework for native plant restoration in the Mojave Desert. *Ecology and Evolution* 12(4) April 2022: 1-16. <https://onlinelibrary.wiley.com/doi/pdfdirect/10.1002/ece3.8805>

Webb, R.H. 2002. Recovery of Severely Compacted Soils in the Mojave Desert, California, USA.
Arid Land Research and Management 16: 291-305.